

MINI REVIEW

O-GlcNAcylation of key nuclear and cytoskeletal proteins: reciprocity with O-phosphorylation and putative roles in protein multimerization

Gerald W.Hart¹, Lisa K.Kreppel, Frank I.Comer, C.Shane Arnold, Doris M.Snow, Zhengyi Ye, Xiaogang Cheng, Debbie DellaManna, Drury S.Caine, Betty J.Earles, Yoshihiro Akimoto, Robert N.Cole and Bradley K.Hayes

Department of Biochemistry and Molecular Genetics, University of Alabama at Birmingham, School of Medicine and Dentistry, 1918 University Boulevard, Birmingham, AL 35294-0005, USA

¹To whom correspondence should be addressed

Key words: O-GlcNAc/phosphorylation/transcription/glycosyltransferase/regulation

Introduction

O-Linked N-acetylglucosamine, O-GlcNAc, was discovered (Torres and Hart, 1984) as part of our studies using highly-purified glycosyltransferases to investigate saccharide structures on lymphocytes and tumor cells (Powell *et al.*, 1987; Whiteheart and Hart, 1987; Passaniti and Hart, 1988; Reichner *et al.*, 1988; Whiteheart *et al.*, 1989). Our glycosyltransferase studies are a direct outgrowth of the pioneering work of Professor Hill and his colleagues, which developed the methods for the purification of glycosyltransferases (Paulson *et al.*, 1977; Beyer *et al.*, 1980, 1981; Sadler *et al.*, 1979, 1981, 1982). In particular, it was the purification (Trayer and Hill, 1971; Barker *et al.*, 1972) and ready availability of bovine milk galactosyltransferase that allowed us to employ the enzyme as a highly specific and sensitive probe for O-GlcNAc on nuclear and cytoplasmic proteins (for methods, see Roquemore *et al.*, 1994b). Using this enzymatic probe and other methods, we and others (for review, Hart *et al.*, 1989, 1995a–c) have now documented that O-GlcNAc is ubiquitous and abundant on nuclear and cytoskeletal proteins of virtually all eukaryotes, including protozoans (Ortega-Barria *et al.*, 1990; Dieckmann-Schuppert *et al.*, 1993, 1994; Stanley *et al.*, 1995) and fungi (Machida and Jigami, 1994). In addition, O-GlcNAc is highly dynamic, with turnover rates much higher than the protein backbones to which it is attached (Kearse and Hart, 1991; Chou *et al.*, 1992; Roquemore *et al.*, 1996). Virtually all O-GlcNAc proteins examined to date are also phosphoproteins, and in some instances Ser(Thr)-O-GlcNAc and Ser(Thr)-O-phosphate appear to reciprocally occupy the same hydroxyl groups (Kelly *et al.*, 1993; Chou *et al.*, 1995a,b). The dynamic characteristics of O-GlcNAc, and the importance of O-GlcNAcylated proteins in cellular regulation and in cytoskeletal structure, have led us to hypothesize that O-GlcNAc is a regulatory modification that not only regulates protein phosphorylation, but also is involved in modulating protein multimerization (Hart *et al.*, 1995b,c).

O-GlcNAcylation is most abundant in the nucleus

Early studies using highly-purified rat liver subcellular fractions (Holt and Hart, 1986), demonstrated that most O-GlcNAc bearing proteins are localized to the nucleus or cytoplasm. The highest concentration of O-GlcNAc was found in the nuclear envelope on proteins that appeared to comigrate with nuclear pore proteins. Using monoclonal antibodies to the nuclear pore proteins (Snow *et al.*, 1987), the nucleoporins were found to be multiply O-GlcNAcylated, and the sugar was shown to be a major epitope recognized by nearly all of the monoclonal antibodies (Holt *et al.*, 1987b). Later work has shown that monoclonal antibodies that are selective for O-GlcNAc and the lectin, wheat germ agglutinin (WGA), which binds clustered GlcNAc moieties, block nuclear transport of macromolecules (Wolff *et al.*, 1988; Forbes, 1992; Hanover, 1992). However, capping of many of the O-GlcNAc residues by the *in vitro* attachment of galactose residues does not appear to block nuclear transport of macromolecules (Miller and Hanover, 1994).

Even though the concentration of O-GlcNAc is the highest at the nuclear pore complex (Holt *et al.*, 1987b; Hanover *et al.*, 1987; Schindler *et al.*, 1987), most of the proteins bearing this modification are within the nucleoplasm. Studies of salivary gland polytene chromosomes from *Drosophila melanogaster* dramatically demonstrated high concentrations of O-GlcNAc along the entire length of the chromosome (Kelly and Hart, 1989). Interestingly, FITC-WGA-staining showed an intense banding pattern similar to commonly used chromatin dyes. The concentration of O-GlcNAc proteins is much reduced at puffs, which are active sites of gene transcription (Kelly and Hart, 1989). Purification and galactosyltransferase probing of various chromatin fractions indicated that a myriad of proteins in chromatin bear O-GlcNAc moieties, and that the majority of the O-GlcNAc chromatin proteins are associated with chromatin at least as tightly as are the histones (Kelly and Hart, 1989).

O-GlcNAc is ubiquitous in all eukaryotes examined

Since 1984, our laboratory and several others have identified many of the multitude of proteins that are modified by O-GlcNAc. Figure 1 represents a list of those that have been identified at the time of this writing. The list includes RNA polymerase II (Kelly *et al.*, 1993) and virtually all of its transcriptional regulatory proteins that have been examined (Jackson and Tjian, 1988, 1989; Hart *et al.*, 1989; Lichtsteiner and Schibler, 1989; Reason *et al.*, 1992). In addition, many key cytoskeletal proteins (Luthi *et al.*, 1991; Holt *et al.*, 1987a; King and Hounsell, 1989; Chou *et al.*, 1992; Chou and Omary, 1993; Dong *et al.*, 1993; Ku and Bishr Omary, 1994; Murphy *et al.*, 1994; Ding and Vandre, 1996; Vostal and Krasnewich, 1996), viral proteins (Benko *et al.*, 1988; Mullis *et al.*, 1990;

O-GlcNAcylated Proteins

- Transcription - Pol II & Reg. Factors.
- Cytoskeleton - Intermediate Filaments; MAP Proteins, 'Bridging Proteins'.
- Tumor Suppressors & Oncogenes.
- Viral Proteins - Tegument & Fiber.
- Phosphatases & Kinases.
- Heat Shock Proteins.
- Nuclear Pore Proteins

Fig. 1. Identified O-GlcNAcylated proteins. Only a very small percentage of the total O-GlcNAcylated proteins have been identified to date. For references, see text

Gonzalez and Burrone, 1992; Whitford and Faulkner, 1992; Greis *et al.*, 1994), nuclear oncogenes (Privalsky, 1990; Chou *et al.*, 1995a,b), tumor suppressors (Shaw *et al.*, 1996), fungal DNA-binding proteins (Machida and Jigami, 1994), steroid receptors (Jiang *et al.*, unpublished observations), small heat shock proteins (Roquemoire *et al.*, 1992; Koga *et al.*, 1996), phosphatases (Meikrantz *et al.*, 1991), kinases (Matsuoka and Hart, unpublished observations), and kinase regulatory proteins (Datta *et al.*, 1989) have been found to be modified by O-GlcNAc. The common feature of these O-GlcNAcylated proteins is that they are all also phosphoproteins and most are known to reversibly form multimeric associations depending upon their phosphorylation states.

O-GlcNAc is a regulatory modification that can be dynamically reciprocal with phosphorylation

The location of O-GlcNAc at Ser(Thr) sites that are similar or identical to those used by many kinases, and the presence of the modification on many of the cell's most important proteins led us to speculate that O-GlcNAc might be a regulatory modification analogous to phosphorylation. Studies of the activation of murine lymphocytes by mitogens and antigens demonstrated that O-GlcNAc levels on lymphocyte proteins change within minutes after stimulation of the cells (Kearse and Hart, 1991). Pulse-chase studies on cytokeratins (Chou *et al.*, 1992) and on the small heat shock proteins (Roquemoire *et al.*, 1996) clearly show that O-GlcNAc turns over many times faster than the polypeptide to which it is attached. These studies, and those that show the reciprocal nature of Ser(Thr)-O-phosphorylation and Ser(Thr)-O-GlcNAcylation (Kelly *et al.*, 1993; Chou *et al.*, 1995a), led us to suggest that the posttranslational modification of many nuclear and cytoplasmic proteins is a ternary process *not* a binary one (Figure 2). Thus, the modulation of some protein's functions may be more complicated than the simple addition and removal of phosphate. Rather, it is clear that some proteins exist in up to three different isoforms, naked polypeptide, phosphorylated polypeptide and O-GlcNAcylated polypeptide, each with distinct physical and biological properties. Thus far, circumstantial evidence suggests the working hypoth-

esis that O-GlcNAcylation of proteins generally favors their associations with other proteins, whereas phosphorylation often appears to favor disassociation. We have evaluated the effect of blocking O-GlcNAc turnover in live cells by transfecting them with cDNAs that encode cytoplasmic or nuclear localizing forms of bovine milk galactosyltransferase, an enzyme that 'caps' terminal GlcNAc residues by transferring galactose from UDP-galactose to form Gal β 1-4GlcNAc linkages (Snow *et al.*, 1996). The transfection of Chinese hamster ovary cells with these cDNAs appears to kill the cells within one cell cycle, whereas transfection with the cDNA encoding the normal luminal Golgi localizing enzyme has no effect on cell viability. These data are consistent with the hypothesis that the dynamic aspects of O-GlcNAcylation are important to the life of the cell. However, the mechanisms by which the nuclear or cytoplasmically localized galactosyltransferase kills the cell remains undetermined.

O-GlcNAcylation sites are similar or identical to kinase sites

As part of our effort to determine the biological functions of O-GlcNAcylation, we and other laboratories have mapped the sites of attachment of the saccharide on many different proteins (Hart *et al.*, 1995b). The overall approach for site mapping of proteins, as illustrated for the c-myc protein, is shown in Figure 3. While about half of the attachment sites for O-GlcNAc occur at or near a 'PVS' type motif, about half of the sites have no obvious 'consensus' sequence (Hart *et al.*, 1995b). Many of these sites are similar to those used by kinases, particularly glycogen synthase kinase 3 (Ishiguro *et al.*, 1993; Skurat and Roach, 1996), and the MAP kinases such as erk2 kinase (Boulton *et al.*, 1991). Site mapping is allowing site-directed mutagenesis approaches to be carried-out in order to evaluate functions. However, if mutation of a specific serine or threonine residue affects a protein functions, it may be difficult to determine if the loss of function is due to inability to attach O-GlcNAc or to attach O-phosphate. Recently, the major site of




Fig. 2. Dynamic model of the relationship of Ser(Thr)-O-GlcNAc and Ser(Thr)-O-phosphate. In this model, there is often a reciprocal relationship between phosphorylation of a hydroxyl moiety and its O-GlcNAcylation. However, it is anticipated that attachment of O-GlcNAc to nearby or adjacent serine or threonine residues may also influence their phosphorylation.




Fig. 3. General scheme for locating the attachment sites of O-GlcNAc. Trypsin has proven to be most useful for fragmentation. It is helpful to 'tag' the glycopeptides using UDP-[³H]galactose and galactosyltransferase (GT). Generally, several runs of reverse-phase high performance chromatography (RP-HPLC) under different conditions are required. Site location requires a combination of techniques including, gas phase microsequencing, solid-phase manual Edman degradation, and mass spectrometry.

O-GlcNAcylation of the c-myc oncogene protein was mapped to the transactivation domain (Chou *et al.*, 1995b) at Thr-58 (Chou *et al.*, 1995a), which is also a major phosphorylation site and the most important 'hotspot' in mutations in c-myc associated with lymphomas. Mutation of Thr-58 in c-myc to alanine or to methionine is associated with increased tumorigenicity, suggesting that posttranslational modification at Thr-58 leads to negative regulation of growth and neoplastic phenotype. Clearly, given the reciprocal occupancy of O-GlcNAc and phosphate at this hydroxyl, it is now unclear which of these modifications are beneficial in preventing the oncogenic properties of this protein. However, based upon the literature discussing the role of phosphorylation in the binding of tumor suppressor proteins to oncogenes (Henriksson *et al.*, 1993; Pulverer *et al.*, 1994), it appears likely that the form of c-myc that interacts with tumor suppressors like the retinoblastoma (Rb) protein (Lees *et al.*, 1991; Picksley and Lane, 1994; Weinberg, 1995) or p53 protein (White, 1994; Kastan *et al.*, 1995), is the O-GlcNAcylated form. Recently, the p53 tumor suppressor was shown to bear O-GlcNAc (Shaw *et al.*, 1996). These studies suggested that the O-GlcNAcylated form of p53 has higher biological activity and higher DNA binding activity than the non-O-GlcNAcylated species of p53. Recent studies have also shown that SV40 Large T antigen is O-GlcNAcylated (Medina-Vera and Haltiwanger, 1994). Several O-GlcNAcylated DNA-binding proteins have also been described in the filamentous fungus, *Aspergillus oryzae*, suggesting that O-GlcNAcylation may be important in the functions of transcription factors from even the most primitive eukaryotes (Machida and Jigami, 1994).

O-GlcNAc is involved in transcriptional initiation

The 'mucin-like' sequence of the C-terminal domain (CTD) of RNA polymerase II (Pol II) prompted us to investigate if this protein is O-GlcNAcylated (Kelly *et al.*, 1993). All eukaryotic

Pol II proteins have multiple repeats of the consensus sequence, -Tyr-Ser-Pro-Thr-Ser-Pro-Ser-, at their C-terminus (Corden, 1990). Available data indicate that the CTD is unphosphorylated in the transcriptional preinitiation complex (Lu *et al.*, 1991), but becomes heavily phosphorylated with the addition of as many as fifty phosphates preceding transcriptional elongation. We found that a subset of Pol II CTDs are heavily O-GlcNAcylated but contain no phosphates (Kelly *et al.*, 1993). These data suggest a model in which the O-GlcNAcylated isoform of Pol II assembles in the preinitiation complex. The O-GlcNAcs then are a rapidly removed prior to the phosphorylation of these same sites. Phosphorylation of the CTD may be important for breaking apart the initiation complex and allows elongation to take place (Zawel and Reinberg, 1995). This model of the role of O-GlcNAc in transcriptional initiation is consistent with the proposed role of O-GlcNAc in mediating subunit interactions and with its observed reciprocal relationship with phosphorylation. We further postulate that site-specific occupancy of transcription factors and Pol II by bulky O-GlcNAc residues might play an important role in regulating cell specific assembly of the transcriptional initiation complex and thus directly affect cell specific gene expression. Very recently, strong support for a role of O-GlcNAc in transcriptional initiation has been obtained by the ability of specific O-GlcNAcase inhibitors and O-GlcNAc-modified CTD peptides, but not O-GalNAc modified CTD peptides to block mRNA transcription mediated by the major late promoter of adenovirus (Comer and Hart, 1996). Elucidation of the nature of O-GlcNAc's role in transcription will require careful kinetic and biochemical analyses of the relationships of O-GlcNAcylation and phosphorylation on each of the many O-GlcNAc bearing proteins comprising the transcriptional machinery.

O-GlcNAc may mediate cytoskeletal assembly and organization

Site mapping studies on neurofilament proteins indicated that the different types of neurofilaments, H, M, or L, are O-GlcNAcylated (Dong *et al.*, 1993). Based upon prior site-directed mutagenesis studies of these neurofilaments (Gill *et al.*, 1990), the locations of the O-GlcNAc moieties suggest that the saccharide is directly involved in mediating neurofilament assembly. In fact, most of the O-GlcNAc residues on neurofilaments are buried and are inaccessible to galactosyltransferase or hexosaminidase treatments, unless the protein is first denatured or fragmented with protease treatment. The O-GlcNAc moieties on α -crystallins (Roquemore *et al.*, 1992), and those on the adenovirus fiber proteins (Mullis *et al.*, 1990) are also buried on the native molecules. During the analyses of partially purified neurofilaments from rat spinal cord (Dong *et al.*, 1993), we observed an apparently heavily O-GlcNAcylated protein that appeared to be the microtubule-associated protein, tau (Mandelkow and Mandelkow, 1994; Goedert *et al.*, 1995; Ingram, 1995). In normal neurons, tau plays a key role in organizing microtubules in the axons. However, in neurons from Alzheimer disease patients, tau is abnormally hyperphosphorylated causing it to form abnormal filaments (PHF-tau), which may be involved in neuronal death (Mandelkow and Mandelkow, 1993; Mandelkow *et al.*, 1993). We have shown that normal tau from bovine brain is heavily O-GlcNAcylated, with apparently over 12 O-GlcNAcylation sites and an average stoichiometry of ~4 O-GlcNAc residues per mole of protein

(Arnold *et al.*, 1994). These studies suggest that the abnormal hyperphosphorylation of tau in Alzheimer disease neurons may result from a defect in the normal O-GlcNAcylation of serine or threonine residues that allows abnormal phosphorylation to take place. A putative role for O-GlcNAcylation in Alzheimer disease is even more intriguing with the recent finding of O-GlcNAc on the β -amyloid precursor protein (Griffith *et al.*, 1995), and the direct demonstration of altered O-GlcNAc levels in Alzheimer patients (Griffith and Schmitz, 1995). Likewise, a possible role for O-GlcNAc in generally mediating the associations of microtubule binding proteins must be considered in the light of recent reports describing the O-GlcNAcylation of the high molecular weight MAP proteins, including MAP2 (Ding and Vandre, 1996). It is noteworthy that several of the O-GlcNAcylated cytoskeletal proteins, such as band 4.1 (Holt *et al.*, 1987a), vinculin (Vostal and Krasnewich, 1996), talin (Hagmann *et al.*, 1992), synapsin (Luthi *et al.*, 1991), and others are involved in the phosphorylation-dependent reversible bridging of the cytoskeleton to the membrane or other structures. Thus, while direct data are still lacking, it is our working hypothesis that O-GlcNAcylation plays an important role in mediating protein-protein interactions involved in a wide variety of cellular functions, including the organization of the cytoskeleton.

Nuclear and cytoplasmic enzymes that remove or attach O-GlcNAc to proteins are abundant

Analogous to phosphatases, we have identified and purified a cytoplasmic O-GlcNAcase from rat liver that has selectivity for O-GlcNAc peptides (Dong and Hart, 1994). O-GlcNAcase activities and nonspecific hexosaminidases are widely abundant in all cell types we have examined, making the quantification of O-GlcNAc stoichiometry problematic (see Roquemore *et al.*, 1992, 1994a, for discussion). The O-GlcNAcase we have described and purified is similar to a partially purified class of enzymes (Minami *et al.*, 1981; Overdijk *et al.*, 1981; Ueno and Yuan, 1991; Koga *et al.*, 1996) referred to as hexosaminidase C. As yet we know almost nothing about the regulation of O-GlcNAcases, but the cloning of the cDNA encoding this rat cytosolic enzyme is in process.

Using synthetic peptide substrates we have also purified a UDP-GlcNAc:polypeptide O-N-acetylglucosaminyltransferase (O-GlcNAc transferase) from rat liver (Haltiwanger *et al.*, 1990, 1992). The rat liver enzyme appears to have two immunologically related subunits, of Mr = 110,000, and Mr = 78,000 with a molar ratio of 2:1, respectively, in the native enzyme. Photolabeling studies indicate that the 110K subunit contains the active site of the enzyme. The enzyme also appears to be phosphorylated on tyrosine residues and is itself O-GlcNAcylated (Kreppel and Hart, 1996). Using sequence information from 11 tryptic peptides of the catalytic subunit, we have recently cloned the O-GlcNAc transferase from rat liver. The protein is unlike any known glycosyltransferase previously described (Paulson and Colley, 1989; Joziassse, 1992) and is very highly conserved at the primary sequence level based upon a putative open reading frame in the GenBank database obtained from *C.elegans*. Identification of clones encoding the O-GlcNAc transferase has been confirmed by the comparison of the sequences of numerous peptides, overexpression of the protein and corresponding enzyme activity in heterologous systems, and by direct immunoprecipitation and/or immunodepletion of O-GlcNAc transferase activity using a

polyclonal antibody made against the recombinant protein overexpressed in *E.coli*. The availability of the cloned DNA encoding an O-GlcNAc transferase should allow rapid progress on the evaluation of both the regulation of O-GlcNAcylation and the determinations of the saccharide's functions in a wide variety of biological systems.

Acknowledgements

This review is dedicated to Professor Robert L.Hill, Ph.D., Biochemistry Department, Duke University Medical Center, on the occasion of his 65th birthday, and it is based upon a seminar presented at the 'Hillfest' Symposium on the Structure, Function and Evolution of Glycoproteins and Related Molecules, which took place at Orcas Island, Eastsound, Washington from August 16–19, 1995. This work was supported by NIH grants to G.W.H. from R01 HD-13563, R01 CA-42486, a grant from the Juvenile Diabetes Foundation and the Mizutani Foundation, and a gift from Oxford Glycosystems, Inc. F.I.C. is a predoctoral fellow of the National Science Foundation.

References

- Arnold, C.S., Cole, R.N., Johnson, G.V.W. and Hart, G.W. (1994) Tau is a Glycoprotein. *Soc. Neurosci. Abstr.*, **20**, 1035.
- Barker, R., Olsen, K.W., Shaper, J.H. and Hill, R.L. (1972) Agarose derivatives of uridine diphosphate and N-acetylglucosamine for the purification of a galactosyltransferase. *J. Biol. Chem.*, **247**, 7135–7147.
- Benko, D.M., Haltiwanger, R.S., Hart, G.W. and Gibson, W. (1988) Virion basic phosphoprotein from human cytomegalovirus contains O-linked N-acetylglucosamine. *Proc. Natl. Acad. Sci. USA*, **85**, 2573–2577.
- Beyer, T.A., Sadler, J.E. and Hill, R.L. (1980) Purification to homogeneity of H blood group beta-galactoside alpha 1-2 fucosyltransferase from porcine submaxillary gland. *J. Biol. Chem.*, **255**, 5364–5372.
- Beyer, T.A., Sadler, J.E., Rearick, J.I., Paulson, J.C., and Hill, R.L. (1981) Glycosyltransferases and their use in assessing oligosaccharide structure and structure-function relationships. *Adv. Enzymol.*, **52**, 23–175.
- Boulton, T.G., Nye, S.H., Robbins, D.J., Ip, N.Y., Radziejewska, E., Morgenbesser, S.D., DePinho, R.A., Panayotatos, N., Cobb, M.H. and Yancopoulos, G.D. (1991) ERKs: a family of protein-serine/threonine kinases that are activated and tyrosine phosphorylated in response to insulin and NGF. *Cell*, **65**, 663–675.
- Chou, C.-F., Smith, A.J. and Omary, M.B. (1992) Characterization and dynamics of O-linked glycosylation of human cytokeratin 8 and 18. *J. Biol. Chem.*, **267**, 3901–3906.
- Chou, C.-F. and Omary, M.B. (1993) Mitotic arrest-associated enhancement of O-linked glycosylation and phosphorylation of human keratins 8 and 18. *J. Biol. Chem.*, **268**, 4465–4472.
- Chou, T.-Y., Hart, G.W. and Dang, C.V. (1995a) c-Myc is glycosylated at threonine 58, a known phosphorylation site and a mutational hot spot in lymphomas. *J. Biol. Chem.*, **270**, 18961–18965.
- Chou, T.-Y., Dang, C.V. and Hart, G.W. (1995b) Glycosylation of the c-myc transactivation domain. *Proc. Natl. Acad. Sci. USA*, **92**, 4417–4421.
- Comer, F.I. and Hart, G.W. (1996) Investigating the role of O-GlcNAc on RNA polymerase II. *FASEB J.*, **10**, A1119.
- Corden, J.L. (1990) Tails of RNA polymerase II. *Trends Biol. Sci.*, **15**, 383–387.
- Datta, B., Ray, M.K., Chakrabarti, D., Wylie, D.E. and Gupta, N.K. (1989) Glycosylation of eukaryotic peptide chain initiation factor 2 (eIF-2)-associated 67-kDa polypeptide (p⁶⁷) and its possible role in the inhibition of eIF-2 kinase-catalyzed phosphorylation of the eIF-2 α -subunit. *J. Biol. Chem.*, **264**, 20620–20624.
- Dieckmann-Schuppert, A., Bause, E. and Schwarz, R.T. (1993) Studies on O-glycans of *Plasmodium falciparum*-infected human erythrocytes—evidence for O-GlcNAc and O-GlcNAc-transferase in malaria parasites. *Eur. J. Biochem.*, **216**, 779–788.
- Dieckmann-Schuppert, A., Bause, E. and Schwarz, R.T. (1994) Glycosylation reactions in *Plasmodium falciparum*, *Toxoplasma gondii*, and *Trypanosoma brucei* probed by the use of synthetic peptides. *Biochim. Biophys. Acta Gen. Subj.*, **1199**, 37–44.
- Ding, M. and Vandre, D.D. (1996) High molecular weight microtubule-associated proteins contain O-linked N-acetylglucosamine. *J. Biol. Chem.*, **271**, 12555–12561.
- Dong, D.L.-Y., Xu, Z.-S., Chevrier, M.R., Cotter, R.J., Cleveland, D.W. and Hart, G.W. (1993) Glycosylation of mammalian neurofilaments. Localiza-

- tion of multiple O-linked N-acetylglucosamine moieties on neurofilament polypeptides L and M. *J. Biol. Chem.*, **268**, 16679–16687.
- Dong, D.L.-Y. and Hart, G.W. (1994) Purification and characterization of an O-GlcNAc selective N-acetyl- β -D-glucosaminidase from rat spleen cytosol. *J. Biol. Chem.*, **269**, 19321–19330.
- Dong, L.-Y.D., Xu, Z.-S., Chevrier, M.R., Cotter, R.J., Cleveland, D.W. and Hart, G.W. (1993) Glycosylation of mammalian neurofilaments. Localization of multiple O-linked N-acetylglucosamine moieties on neurofilament polypeptides L and M. *J. Biol. Chem.*, **268**, 16679–16687.
- Forbes, D.J. (1992) Structure and function of the nuclear pore complex. *Annu. Rev. Cell Biol.*, **8**, 495–527.
- Gill, S.R., Wong, P.C., Monteiro, M.J. and Cleveland, D.W. (1990) Assembly properties of dominant and recessive mutations in the small mouse neurofilament (NF-L) subunit. *J. Cell Biol.*, **111**, 2005–2019.
- Goedert, M., Jakes, R., Spillantini, M.G., Crowther, R.A., Cohen, P., Vanmechelen, E., Probst, A., Göttsch, J. and Bürki, K. (1995) Tau protein in Alzheimer's disease. *Biochem. Soc. Trans.*, **23**, 80–86.
- Gonzalez, S.A. and Burrone, O.R. (1992) Rotavirus NS26 is modified by addition of single O-linked residues of N-acetylglucosamine. *Virology*, **182**, 8–16.
- Greis, K., Gibson, W. and Hart, G.W. (1994) Site-specific glycosylation of the human cytomegalovirus tegument basic phosphoprotein (UL32) at serine 921 and serine 952. *J. Virol.*, **68**, 9339–9349.
- Griffith, L.S. and Schmitz, B. (1995) O-Linked N-acetylglucosamine is upregulated in Alzheimer brains. *Biochem. Biophys. Res. Commun.*, **213**, 424–431.
- Griffith, L.S., Mathes, M. and Schmitz, B. (1995) b-Amyloid precursor protein is modified with O-linked N-acetylglucosamine. *J. Neurosci. Res.*, **41**, 270–278.
- Hagmann, J., Grob, M. and Burger, M.M. (1992) The cytoskeletal protein talin is O-glycosylated. *J. Biol. Chem.*, **267**, 14424–14428.
- Haltiwanger, R.S., Holt, G.D. and Hart, G.W. (1990) Enzymatic addition of O-GlcNAc to nuclear and cytoplasmic proteins. Identification of a uridine diphospho-N-acetylglucosamine:peptide β -N-acetylglucosaminyltransferase. *J. Biol. Chem.*, **265**, 2563–2568.
- Haltiwanger, R.S., Blomberg, M.A. and Hart, G.W. (1992) Glycosylation of nuclear and cytoplasmic proteins. Purification and characterization of a uridine diphospho-N-acetylglucosamine:polypeptide β -N-acetylglucosaminyltransferase. *J. Biol. Chem.*, **267**, 9005–9013.
- Hanover, J.A. (1992) The nuclear pore: at the crossroads. *FASEB J.*, **6**, 2288–2295.
- Hanover, J.A., Cohen, C.K., Willingham, M.C. and Park, M.K. (1987) O-Linked N-acetylglucosamine is attached to proteins of the nuclear pore. Evidence for cytoplasmic and nucleoplasmic glycoproteins. *J. Biol. Chem.*, **262**, 9887–9894.
- Hart, G.W., Haltiwanger, R.S., Holt, G.D. and Kelly, W.G. (1989) Glycosylation in the nucleus and cytoplasm. *Annu. Rev. Biochem.*, **58**, 841–874.
- Hart, G.W., Chou, T.-Y., Jiang, M.-S., Greis, K.D., Cole, R.N., Comer, F.I., Arnold, C.S., Matsuoka, T., Snow, D.M., Hayes, B.K., Kreppel, L.K. and Earles, B.J. (1995a) O-GlcNAcylation and phosphorylation reciprocity on nuclear and cytoskeletal proteins. *NIH Catalyst*, **3**, 8–9.
- Hart, G.W., Greis, K.D., Dong, L.-Y.D., Blomberg, M.A., Chou, T.-Y., Jiang, M.-S., Roquemore, E.P., Snow, D.M., Kreppel, L.K., Cole, R.N. and Hayes, B.K. (1995b) Ubiquitous and temporal glycosylation of nuclear and cytoplasmic proteins. *Pure Appl. Chem.*, **67**, 1637–1645.
- Hart, G.W., Greis, K.D., Dong, L.-Y.D., Blomberg, M.A., Chou, T.-Y., Jiang, M.-S., Roquemore, E.P., Snow, D.M., Kreppel, L.K., Cole, R.N., Comer, F.I., Arnold, C.S. and Hayes, B.K. (1995c) O-Linked N-acetylglucosamine: the 'yin-yang' of ser/thr phosphorylation? Nuclear and cytoplasmic glycosylation. *Adv. Exp. Med. Biol.*, **376**, 115–123.
- Henriksson, M., Bakardjiev, A., Klein, G. and Lyscher, B. (1993) Phosphorylation sites mapping in the N-terminal domain of c-myc modulate its transforming potential. *Oncogene*, **8**, 3199–3209.
- Holt, G.D. and Hart, G.W. (1986) The subcellular distribution of terminal N-acetylglucosamine moieties. Localization of a novel protein-saccharide linkage, O-linked GlcNAc. *J. Biol. Chem.*, **261**, 8049–8057.
- Holt, G.D., Haltiwanger, R.S., Torres, C.R. and Hart, G.W. (1987a) Erythrocytes contain cytoplasmic glycoproteins. O-Linked GlcNAc on band 4.1. *J. Biol. Chem.*, **262**, 14847–14850.
- Holt, G.D., Snow, C.M., Senior, A., Haltiwanger, R.S., Gerace, L. and Hart, G.W. (1987b) Nuclear pore complex glycoproteins contain cytoplasmically disposed O-linked N-acetylglucosamine. *J. Cell Biol.*, **104**, 1157–1164.
- Ingram, V.M. (1995) Alzheimer tangles and abnormal phosphorylation. *Science*, **267**, 1889–1890.
- Ishiguro, K., Shiratsuchi, A., Sato, S., Omori, A., Arioka, M., Kobayashi, S., Uchida, T. and Imahori, K. (1993) Glycogen synthase kinase 3b is identical to tau protein kinase I generating several epitopes of paired helical filaments. *FEBS Lett.*, **325**, 167–172.
- Jackson, S.P. and Tjian, R. (1988) O-Glycosylation of eukaryotic transcription factors: Implications for mechanisms of transcriptional regulation. *Cell*, **55**, 125–133.
- Jackson, S.P. and Tjian, R. (1989) Purification and analysis of RNA polymerase II transcription factors by using wheat germ agglutinin affinity chromatography. *Proc. Natl. Acad. Sci. USA*, **86**, 1781–1785.
- Joziasse, D.H. (1992) Mammalian glycosyltransferases: genomic organization and protein structure. *Glycobiology*, **2**, 271–277.
- Kastan, M.B., Canman, C.E. and Leonard, C.J. (1995) P53, cell cycle control and apoptosis: Implications for cancer. *Cancer Metastasis Rev.*, **14**, 3–15.
- Kearse, K.P. and Hart, G.W. (1991) Lymphocyte activation induces rapid changes in nuclear and cytoplasmic glycoproteins. *Proc. Natl. Acad. Sci. USA*, **88**, 1701–1705.
- Kelly, W.G. and Hart, G.W. (1989) Glycosylation of chromosomal proteins: localization of O-linked N-acetylglucosamine in *Drosophila* chromatin. *Cell*, **57**, 243–251.
- Kelly, W.G., Dahmus, M.E. and Hart, G.W. (1993) RNA polymerase II is a glycoprotein. Modification of the COOH-terminal domain by O-GlcNAc. *J. Biol. Chem.*, **268**, 10416–10424.
- King, I.A. and Hounsell, E.F. (1989) Cytokeratin 13 contains O-glycosidically linked N-acetylglucosamine residues. *J. Biol. Chem.*, **264**, 14022–14028.
- Koga, D., Hoshika, H., Matsushita, M., Tanaka, A., Ide, A. and Kono, M. (1996) Purification and characterization of beta-N-acetylhexosaminidase from the liver of a prawn, *Penaeus japonicus*. *Biosci. Biotechnol. Biochem.*, **60**, 194–199.
- Kreppel, L.K. and Hart, G.W. (1996) Cloning and characterization of O-GlcNAc transferase: a novel highly conserved enzyme. *FASEB J.*, **10**, A1120.
- Ku, N.-O. and Bishr Omary, M. (1994) Expression, glycosylation, and phosphorylation of human keratins 8 and 18 in insect cells. *Exp. Cell Res.*, **211**, 24–35.
- Lees, J.A., Buchkovich, K.J., Marshak, D.R., Anderson, C.W. and Harlow, E. (1991) The retinoblastoma protein is phosphorylated on multiple sites by human cdc2. *EMBO J.*, **10**, 4279–4290.
- Lichtsteiner, S. and Schibler, U. (1989) A glycosylated liver-specific transcription factor stimulates transcription of the albumin gene. *Cell*, **57**, 1179–1187.
- Lu, H., Flores, O., Weinmann, R. and Reinberg, D. (1991) The nonphosphorylated form of RNA polymerase II preferentially associates with the preinitiation complex. *Proc. Natl. Acad. Sci. USA*, **88**, 10004–10008.
- Luthi, T., Haltiwanger, R.S., Greengard, P. and Bahler, M. (1991) Synapsins contain O-linked N-acetylglucosamine. *J. Neurochem.*, **56**, 1493–1498.
- Machida, M. and Jigami, Y. (1994) Glycosylated DNA-binding proteins from filamentous fungus, *Aspergillus oryzae*: modification with N-acetylglucosamine monosaccharide through an O-glycosidic linkage. *Biosci. Biotechnol. Biochem.*, **58**, 344–348.
- Mandelkow, E. and Mandelkow, E.-M. (1994) Microtubule structure. *Curr. Opin. Struct. Biol.*, **4**, 171–179.
- Mandelkow, E.-M. and Mandelkow, E. (1993) Tau as a marker for Alzheimer's disease. *Trends Biol. Sci.*, **18**, 480–483.
- Mandelkow, E.-M., Biernat, J., Drewes, G., Steiner, B., Lichtenberg-Kraag, B., Wille, H., Gustke, N. and Mandelkow, E. (1993) Microtubule-associated protein tau, paired helical filaments, and phosphorylation. *Ann. NY Acad. Sci.*, **695**, 209–216.
- Medina-Vera, L. and Haltiwanger, R.S. (1994) SV40 large T antigen is modified with O-linked N-acetylglucosamine. *Mol. Biol. Cell*, **5**, 340a.
- Meikrantz, W., Smith, D.M., Sladicka, M.M. and Schlegel, R.A. (1991) Nuclear localization of an O-glycosylated protein phosphotyrosine phosphatase from human cells. *J. Cell Sci.*, **98**, 303–307.
- Miller, M.W. and Hanover, J.A. (1994) Functional nuclear pores reconstituted with b1–4 galactose-modified O-linked N-acetylglucosamine glycoproteins. *J. Biol. Chem.*, **269**, 9289–9297.
- Minami, R., Nakamura, F., Oyanagi, K. and Nakao, T. (1981) Hexosaminidase C in brain affected by Tay-Sachs disease. *J. Exp. Med.*, **133**, 175–185.
- Mullis, K.G., Haltiwanger, R.S., Hart, G.W., Marchase, R.B. and Engler, J.A. (1990) Relative accessibility of N-acetylglucosamine in trimers of the adenovirus types 2 and 5 fiber proteins. *J. Virol.*, **64**, 5317–5323.
- Murphy, J.-E., Hanover, J.A., Froehlich, M., DuBois, G. and Keen, J.H. (1994) Clathrin assembly protein AP-3 is phosphorylated and glycosylated on the 50-kDa structural domain. *J. Biol. Chem.*, **269**, 21346–21352.
- Ortega-Barria, E., Ward, H.D., Evans, J.E. and Pereira, M.E.A. (1990) N-Acetyl-D-glucosamine is present in cysts and trophozoites of *Giardia lamblia* and serves as receptor for wheatgerm agglutinin. *Mol. Biochem. Parasitol.*, **43**, 151–166.

- Overdijk, B., Van Der Kroef, W.M.J., Van Steijn, G.E.J. and Lisman, J.J.W. (1981) Isolation and further characterization of bovine brain hexosaminidase C. *Biochim. Biophys. Acta*, **659**, 255–266.
- Passaniti, A. and Hart, G.W. (1988) Cell surface sialylation and tumor metastasis: metastatic potential of B16 melanoma variants correlates with their relative numbers of specific penultimate oligosaccharide structures. *J. Biol. Chem.*, **263**, 7591–7603.
- Paulson, J.C. and Colley, K.J. (1989) Glycosyltransferases. Structure, localization, and control of cell type-specific glycosylation. *J. Biol. Chem.*, **264**, 17615–17618.
- Paulson, J.C., Beranek, W.E. and Hill, R.L. (1977) Purification of a sialyltransferase from bovine colostrum by affinity chromatography on CDP-agarose. *J. Biol. Chem.*, **252**, 2356–2362.
- Picksley, S.M. and Lane, D.P. (1994) p53 and Rb: their cellular roles. *Curr. Opin. Cell Biol.*, **6**, 853–858.
- Powell, L.D., Whiteheart, S.W. and Hart, G.W. (1987) Cell surface sialic acid influences tumor cell recognition in the mixed lymphocyte reaction. *J. Immunol.*, **139**, 262–270.
- Privalsky, M.L. (1990) A subpopulation of the avian erythroblastosis virus v-erbA protein, a member of the nuclear hormone receptor family, is glycosylated. *J. Virol.*, **64**, 463–466.
- Pulverer, B.J., Fisher, C., Voudsen, K., Littlewood, T., Evan, G. and Woodgett, J.R. (1994) Site-specific modulation of c-Myc cotransformation by residues phosphorylated *in vivo*. *Oncogene*, **9**, 59–70.
- Reason, A.J., Morris, H.R., Panico, M., Marais, R., Treisman, R.H., Haltiwanger, R.S., Hart, G.W., Kelly, W.G. and Dell, A. (1992) Localization of O-GlcNAc modification on the serum response transcription factor. *J. Biol. Chem.*, **267**, 16911–16921.
- Reichner, J.S., Whiteheart, S.W. and Hart, G.W. (1988) Intracellular trafficking of cell surface sialoglycoconjugates. *J. Biol. Chem.*, **263**, 16316–16326.
- Roquemore, E.P., Dell, A., Morris, H.R., Panico, M., Reason, A.J., Savoy, L.-A., Wistow, G.J., Zigler, J.S., Jr., Earles, B.J. and Hart, G.W. (1992) Vertebrate lens α -crystallins are modified by O-linked N-acetylglucosamine. *J. Biol. Chem.*, **267**, 555–563.
- Roquemore, E.P., Chou, T.-Y. and Hart, G.W. (1994a) Detection of O-linked N-acetylglucosamine (O-GlcNAc) on cytoplasmic and nuclear proteins. *Methods Enzymol.*, **230**, 443–460.
- Roquemore, E.P., Chou, T.-Y. and Hart, G.W. (1994b) Detection of O-linked N-acetylglucosamine (O-GlcNAc) on cytoplasmic and nuclear proteins. *Methods Enzymol.*, **230**, 443–460.
- Roquemore, E.P., Chevrier, M.R., Cotter, R.J. and Hart, G.W. (1996) Dynamic O-GlcNAcylation of the small heat shock protein α B-crystallin. *Biochemistry*, **35**, 3578–3586.
- Sadler, J.E., Rearick, J.I., Paulson, J.C. and Hill, R.L. (1979) Purification to homogeneity of a beta-galactoside α 2 leads to 3 sialyltransferase and partial purification of an alpha-N-acetylgalactosaminide α 2 leads to 6 sialyltransferase from porcine submaxillary glands. *J. Biol. Chem.*, **254**, 4434–4442.
- Sadler, J.E., Beyer, T.A. and Hill, R.L. (1981) Affinity chromatography of glycosyltransferases. *J. Chromatogr.*, **215**, 181–194.
- Sadler, J.E., Beyer, T.A., Oppenheimer, C.L., Paulson, J.C., Prieels, J.P., Rearick, J.I. and Hill, R.L. (1982) Purification of mammalian glycosyltransferases. *Methods Enzymol.*, **83**, 458–514.
- Schindler, M., Hogan, M., Miller, R. and DeGaetano, D. (1987) A nuclear specific glycoprotein representative of a unique pattern of glycosylation. *J. Biol. Chem.*, **262**, 1254–1260.
- Shaw, P., Freeman, J., Bovey, R. and Iggo, R. (1996) Regulation of specific DNA binding by p53: evidence for a role for O-glycosylation and charged residues at the carboxy-terminus. *Oncogene*, **12**, 921–930.
- Skurat, A.V. and Roach, P.J. (1996) Multiple mechanisms for the phosphorylation of C-terminal regulatory sites in rabbit muscle glycogen synthase expressed in COS cells. *Biochem. J.*, **313**, 45–50.
- Snow, C.M., Senior, A. and Gerace, L. (1987) Monoclonal antibodies identify a group of nuclear pore complex glycoproteins. *J. Cell Biol.*, **104**, 1143–1156.
- Snow, D.M., Shaper, J.H., Shaper, N.L. and Hart, G.W. (1996) Cytosolic galactosyltransferase mediated capping of O-GlcNAc in CHO cells: a model to elucidate function. *Mol. Biol. Cell*, **6**, 357a.
- Stanley, S.L., Tian, K., Koester, J.P. and Li, E. (1995) The serine-rich *Entamoeba histolytica* protein is a phosphorylated membrane protein containing O-linked terminal N-acetylglucosamine residues. *J. Biol. Chem.*, **270**, 4121–4126.
- Torres, C.-R. and Hart, G.W. (1984) Topography and polypeptide distribution of terminal N-acetylglucosamine residues on the surfaces of intact lymphocytes. *J. Biol. Chem.*, **259**, 3308–3317.
- Trayer, I.P. and Hill, R.L. (1971) The purification and properties of the A protein of lactose synthetase. *J. Biol. Chem.*, **246**, 6666–6675.
- Ueno, R. and Yuan, C.-S. (1991) Purification and properties of neutral b-N-acetylglucosaminidase from carp blood. *Biochim. Biophys. Acta Gen. Subj.*, **1074**, 79–84.
- Vostal, J.G. and Krasnewich, D.M. (1996) Dynamic O-linked N-acetylglucosamine glycosylation of platelet vinculin. *Mol. Biol. Cell*, **5**, 263a.
- Weinberg, R.A. (1995) The retinoblastoma protein and cell cycle control. *Cell*, **81**, 323–330.
- White, E. (1994) Tumour biology: p53, guardian of Rb. *Nature*, **371**, 21–22.
- Whiteheart, S.W., Passaniti, A., Reichner, J.S., Holt, G.D., Haltiwanger, R.S. and Hart, G.W. (1989) Glycosyltransferase probes. *Methods Enzymol.*, **179**, 82–95.
- Whiteheart, S.W. and Hart, G.W. (1987) Sialyltransferases as specific cell surface probes of terminal and penultimate saccharide structures on living cells. *Anal. Biochem.*, **163**, 123–135.
- Whitford, M. and Faulkner, P. (1992) A structural polypeptide of the baculovirus *Autographa californica* nuclear polyhedrosis virus contains O-linked N-acetylglucosamine. *J. Virol.*, **66**, 3324–3329.
- Wolff, B., Willingham, M.C. and Hanover, J.A. (1988) Nuclear protein import: specificity for transport across the nuclear pore. *Exp. Cell Res.*, **178**, 318–334.
- Zawel, L. and Reinberg, D. (1995) Common themes in assembly and function of eukaryotic transcription complexes. *Annu. Rev. Biochem.*, **64**, 533–561.

Received on July 26, 1996; accepted on July 28, 1996